



## Integrated Fertilization to Support Crop Productivity in Saline Soils

Said M. El-Bialy<sup>1</sup>, Taha El-Esawy<sup>1</sup>, Hassan El-Ramady<sup>1\*</sup>, Mohammed E. El-Mahrouk<sup>2</sup>,  
Tamer El-Sakhawy<sup>3</sup>, Alaa El-Dein Omara<sup>3</sup>, Megahed Amer<sup>4</sup>, Mohamed A. El-Serpiny<sup>5</sup>,  
Yahya Eid<sup>6</sup> and Eric C. Brevik<sup>7</sup>



<sup>1</sup> Soil and Water Department, Faculty of Agriculture, Kafrelsheikh University, Kafr El-Sheikh 33516, Egypt

<sup>2</sup> Horticulture Department, Faculty of Agriculture, Kafrelsheikh University, Kafr El-Sheikh 33516, Egypt

<sup>3</sup> Agriculture Microbiology Department, Soil, Water and Environment Research Institute (SWERI), Sakha Agricultural Research Station, Agriculture Research Center (ARC), Kafr El-Sheikh 33717, Egypt

<sup>4</sup> Soils Improvement Dept., Soils, Water and Environment Research Institute (SWERI), Sakha Station, Agricultural Research Center (ARC), Kafr El-Sheikh 33717, Egypt

<sup>5</sup> Soil & Water and Environment Research Institute, Agriculture Research Center, Giza, 12619 Egypt

<sup>6</sup> Poultry Dept., Faculty of Agriculture, Kafrelsheikh University 33516, Kafr El-Sheikh, EGYPT

<sup>7</sup> School of Agricultural Sciences and School of Earth Systems and Sustainability, 1205 Lincoln Drive, Southern Illinois University, Carbondale, IL 62901 USA

**I**NCREASING global population and climate change represent serious challenges to food, water, energy, and soil security for all people. It is thought that climate change will accelerate the harmful impacts of soil salinization and increase the spread of salinity related problems to new areas in the near future. Thus, soil salinity management should be considered a crucial challenge for the future of humanity. This future does not depend only on crop production, but also on other ecosystem services such as the supply of clean water, bioenergy, and related soil products that make modern civilization possible. Soil salinization can reduce crop productivity and other ecosystem services, but proper management, including integrated fertilization, may support crops under such conditions. Integration means utilization of multiple types of fertilizers, including bio-, chemical, and nano fertilizers. Integrated fertilization can mitigate salinity stress by enhancing the biochemical, physiological, and molecular attributes of cultivated plants under salt stress while improving soil properties and conditions. The right combination of fertilizers and cropping systems is still an open question that needs to be answered.

**Keywords:** Mineral fertilizer, Organic fertilizer, Biofertilizers, Nanofertilizers, Oxidative stress.

### 1. Introduction

Crop production depends on many factors, including soil and nutrient conditions. Fertilization can not only boost crop nutrition but also ensure nutrient carryover to future crops (Farias et al. 2025). There are many kinds of fertilizers, including biofertilizers, chemical fertilizers, nano fertilizers, and organic fertilizers. While these fertilizer types can be used independently, they can also be used in combinations designed to take advantage of the best attributes of each, an approach known as integrated fertilization systems (IFS). This approach is also known as integrated nutrient management. Integrated fertilization has many benefits for the soil, crops (including food and forage), and animal production (Simões et al. 2025). Studies confirm that IFS can increase crop productivity and resource use efficiency, along with improving soil quality due to improved soil properties (Wang et al. 2025). Improved soil properties in turn lead to improved soil security and ecosystem services, which supports food, water, and energy security (McBratney et al., 2014; Brevik et al., 2018). Optimizing nutrient and water management is a crucial role of IFS, especially under stressful conditions (Liu et al. 2025). Integrated fertilization can also help address global climate change, as greenhouse gas (GHG) emissions are reduced by ~46% under IFS compared to traditional agriculture (Chen et al. 2025).

\*Corresponding author e-mail: hassan.elramady@agr.kfs.edu.eg

Received: 30/05/2025; Accepted: 11/07/2025

DOI: 10.21608/EJSS.2025.390523.2196

©2025 National Information and Documentation Center (NIDOC)

Soil salinization represents one of the largest global challenges under global climate change. Soil salinization hinders food security and environmental sustainability, and climate change is accelerating the spread of soil salinization into currently unaffected regions (Mukhopadhyay *et al.* 2021). Soil salinity threatens around 10% of

the world's total arable land, with soil salinization due to human activities affecting around 76 million ha (Mishra *et al.* 2023). The reduced crop yields and nutrient imbalances introduced by soil salinity also represent a threat to human health (Brevik, 2010). Therefore, approaches are needed to mitigate and manage soil salinity including (1) soil mapping (Chi *et al.* 2025), (2) physical management such as sub-soiling (Su *et al.* 2025), deep ploughing (Li *et al.* 2025), leveling (El-Ramady *et al.* 2022), sand mixing (Xu *et al.* 2025), leaching (El-Ramady *et al.* 2024), and mulching (Song *et al.* 2025), (3) chemical methods such as gypsum application (Liu Y *et al.* 2025), mineral fertilization (Duan *et al.* 2025), organic fertilization (Hou *et al.* 2025), bio-fertilizers (Dai *et al.* 2025), and nano-fertilizers (Shoukat *et al.* 2024, 2025); (4) remediation through bioremediation (Wang T *et al.* 2025), phytoremediation, and microbial remediation (Zhou *et al.* 2025).

While many papers have been published regarding ways to address soil salinity and crop production, and progress has been made, saline soils still represent a serious challenge to agricultural production. Therefore, this review discusses crop production in saline soils, highlighting the potential role of IFS in mitigating salinity stress.


## 2. Crop production in saline soils

Soil salinity represents a major restriction to global agricultural productivity by imposing a range of physiological, biochemical, and environmental stresses on crops (**Table 1; Figure 1**). Elevated salt concentrations in the soil lead to osmotic stress, which hinders water uptake and creates drought-like conditions even under normal irrigation regime (Majeed and Muhammad, 2019). In addition to water stress, salinity disrupts nutrient homeostasis by interfering with the uptake of essential ions, particularly through sodium-induced inhibition of potassium and calcium absorption (Faisal *et al.*, 2024). This ionic imbalance is further impaired by the generation of reactive oxygen species (ROS), which cause oxidative damage to cellular structures, harm the photosynthetic efficiency, and disrupt key metabolic pathways (Kaviyazhagan and Prakash, 2023). These combined effects led to significant yield reductions, especially in arid and semi-arid regions, thereby increasing food and energy insecurity and increasing the economic burden on farmers through declining crop productivity and rising reclamation costs (Ashford, 2023; Baig *et al.*, 2023). Moreover, the environmental implications of soil salinization contribute to secondary salinization, degradation of soil structure, and loss of microbial and plant biodiversity (Ashford, 2023).

Integrated mitigation strategies have been explored in response to these challenges, including the development of salt-tolerant crop varieties and the implementation of precision irrigation systems (Tarolli *et al.* 2024). However, the success of these strategies depends heavily on a supportive fertilization system designed to address the unique conditions in saline soils. Fertilization plays a pivotal role not only in compensating for nutrient imbalances but also in improving soil physical and biological properties. For example, the use of fertilizers enriched with organic amendments—such as alfalfa residues or pig manure—can enhance soil structure, increase cation exchange capacity, and improve nutrient availability (Liu *et al.* 2024). Additionally, microbial-enriched fertilizers have been shown to stimulate enzyme activity in the soil and promote root development, both of which are critical for enhancing nutrient uptake under salt stress (Wei *et al.* 2024). Nano fertilizers have also been shown to have beneficial effects in salt affected soils (El-Ramady *et al.*, 2024). Innovative systems that integrate nutrient application with salt removal techniques can significantly improve the root zone environment, leading to better crop performance (Wang *et al.* 2024). Equally important is water management, particularly the use of irrigation and drainage systems that facilitate salt leaching and help maintain soil fertility over time (Tarolli *et al.* 2024). It is also important to recognize that plant responses to fertilization under saline conditions are highly variable across species where different crops exhibit unique physiological and biochemical responses due to the complex interactions between salts and nutrients in the rhizosphere (Sheoran 2023).

**Table 1. The response of different crops to soil salinity with and without fertilization programs.**

| Plant species                             | Soil salinity level                                                         | Fertilizer program                                                                                                   | Major study findings                                                                                                                                                              | References                               |
|-------------------------------------------|-----------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------|
| Wheat-maize cropping system               | EC, 3.1-7.8 dS m <sup>-1</sup> during period from 2015 to 2023              | Base fertilizers: 150 kg N /ha, 75 kg P /ha and 50 kg K /ha                                                          | Salinity of irrigation water and soil control salt accumulation in the soil and crop yield                                                                                        | Liu et al. 2024                          |
| Wheat-maize cropping system               | Coastal saline-alkali soil (EC 8.17 dS m <sup>-1</sup> , pH 8.30)           | N, P <sub>2</sub> O <sub>5</sub> , K <sub>2</sub> O: 225, 180 and 100 kg ha <sup>-1</sup>                            | Sodium carboxy-methyl cellulose increased soil water storage and reduced soil salinity                                                                                            | Li et al. 2024                           |
| Sweet sorghum                             | Saline -alkali coastal soil (EC 5.9 dS m <sup>-1</sup> , pH 8.41)           | Total nutrient content (N + P <sub>2</sub> O <sub>5</sub> + K <sub>2</sub> O) was 3, 5 and 2 %, resp.                | Microbial fertilizers boost soil microbial community diversity and network complexity                                                                                             | Wu et al. 2024                           |
| Paddy rice production                     | Soil salinity 2-6 dS m <sup>-1</sup> , pH 9.91                              | Applied N, P <sub>2</sub> O <sub>5</sub> , and K <sub>2</sub> O at 120, 60 and 96 kg·ha <sup>-1</sup> , respectively | Applied biochar increased rice yield by 14.2 %; enhanced N-content in leaves by 28.6 %                                                                                            | Zhao and Grossart (2024)                 |
| Sunflower in pots for 4 months            | Saline-alkali soil (pH 8.77; EC 5.3 dS m <sup>-1</sup> )                    | Without fertilization program                                                                                        | Rhamnolipids increased crop growth by 117%, nutrient availability, and improved soil microbial communities                                                                        | Zhang et al. 2024                        |
| Alfalfa cropping for 3 years              | EC, 12 dS m <sup>-1</sup> top-soil and pH, 10.12                            | Without fertilization program                                                                                        | Cropping alfalfa improves saline soils and soil fertility for sustainable use.                                                                                                    | Zhu et al. 2025                          |
| Cotton in a long-term study (25 years)    | EC, 3.12 – 34.37 dS m <sup>-1</sup>                                         | N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O were 374, 235 and 103 kg·ha <sup>-1</sup> , respectively.      | Soil bacteria helped increase cotton growth; the combination reduced soil salinity                                                                                                | Jia et al. 2025                          |
| Several crops all over the world          | Saline-alkali soils (EC > 4 dS m <sup>-1</sup> or pH > 8.5)                 | Biochar, farmyard manure, and compost on a global level                                                              | Soil EC, pH, and salt content decreased by 23.6, 2.3 and 20.9 % %, compared to without organic amendments                                                                         | Li T et al. 2025                         |
| Tomato plants under greenhouse conditions | Salt stress (60 and 120 mM NaCl; 6-12 dS m <sup>-1</sup> )                  | Applied commercial nutritive solution (Soluponics Universal, Inverfarms, Mexico)                                     | SiO <sub>2</sub> /TiO <sub>2</sub> nano-composites (up to 500 mg L <sup>-1</sup> ) enhanced salt tolerance by stimulate osmolytes production in tomato plants exposed to salinity | Rolón-Cárdenas & Rodríguez-González 2025 |
| Euhalophyte <i>Suaeda salsa</i> L.        | Experimental soil columns (pH 8.48-9.65; EC 0.18 to 70 dS m <sup>-1</sup> ) | Basal fertilizers applied: 10.4 g urea and 7.5 g KH <sub>2</sub> PO <sub>4</sub> to the 0–20 cm soil depth interval  | Plants reduced soil salinity via phyto-desalination; improved soil quality via root-soil interactions                                                                             | Wang et al. 2025                         |
| Paddy and cotton farm lands               | Reclaimed saline-alkali soils for 25 years (EC 11 dS m <sup>-1</sup> )      | Without fertilization program                                                                                        | Bacterial communities important in reclaimed saline-alkaline soil and influenced by pH, SOC, and soil moisture content                                                            | Yin et al. 2025                          |



**Fig. 1. The stresses soil salinity exerts on crops and suggested methods to address them.**

### 3. Integrated fertilization and saline soils

Integrated fertilization is a strategy that combines multiple types of fertilizers (e.g., biofertilizers, chemical fertilizers, nano fertilizers, and organic fertilizers) (**Figure 2**). The IFS approach has proven effective at improving nutrient use efficiency, enhancing soil properties, and mitigating the harmful impacts of soil salinity on crops. This integrated approach supports sustainable crop production under salt-affected soil conditions (Wag *et al.* 2025). Fertilization via industrial chemical fertilizers supplies essential nutrients (e.g., N, P, and K), however, the efficiency of these fertilizers is reduced in salt affected soils due to leaching losses and ion antagonism (Liu *et al.* 2023). Bio-fertilizers such as compost improve the properties of salt-affected soils by increasing soil aggregates and improving water movement through the soil; thus, the salts leach away. Additionally, it supplies the soil and plants with nutrients as the compost decomposes (Hou *et al.* 2025). Nano-fertilizers offer controlled nutrient release in addition to higher surface area per unit mass, enabling precise nutrient delivery and better absorption by higher plant roots (Shoukat *et al.* 2024). Bio-fertilization that includes salt-tolerant microbial strains enhances availability of nutrients in the soil and promotes higher plant stress tolerance. This effect of bio-fertilization improves root architecture and produces growth-promoting substances. Moreover, it leads to better osmotic adjustment (Alotaibi *et al.* 2024). The synergistic influence of combining these different types of fertilization (IFS) helps improve soil structure and nutrient retention as well as maintain soil microbial diversity. **Table 2** illustrates the effect of IFS applied in saline conditions.




Fig. 2. Integrated fertilization against saline conditions.

Table 2. Integrated fertilization to address saline conditions as reported by some recent studies.


| Plant type                          | Salinity conditions                                         | Fertilization type                                                                 | Response                                                                                                                                                                                                                                                        | References                    |
|-------------------------------------|-------------------------------------------------------------|------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|
| Rocket ( <i>Eruca sativa</i> )      | Irrigation by agro-drainage water                           | Nano-fertilizers (NPK), salicylic acid, and poultry wastes                         | The combined treatment of nano-compound fertilizer + poultry wastes + salicylic acid led to improvements in plants, as this approach help the plant resist the adverse effects of saline irrigation water.                                                      | Al-Taey and Al-Musawi, (2019) |
| Wheat ( <i>Triticum aestivum</i> )  | Salt affected soil (5.89 dS $\text{m}^{-1}$ )               | chemical fertilization (NPK) with bio-fertilization on nanoparticles               | This approach is effective as an integrated fertilization system in increasing the values of N, P, and K in grain and straw                                                                                                                                     | Hasan and Saad (2020)         |
| Wheat ( <i>Triticum aestivum</i> )  | Saline-alkali stress (pH 8.13; EC 13.2 dS $\text{m}^{-1}$ ) | Organic amendments combined with bio-stimulants                                    | The highest plant production of enzymatic and non-enzymatic antioxidants was realized when plants were treated with vermicompost and moringa extract                                                                                                            | Awwad <i>et al.</i> (2022)    |
| Wheat ( <i>Triticum aestivum</i> )  | Salt affected soil (EC=7.17dS $\text{m}^{-1}$ )             | Molasses, compost tea, K-humate nano-selenium, nano-manganese, nano-silica         | Organic amendment increased soil cation exchange capacity while nano micronutrients improved wheat grain yield. Co-application of organic amendments and nano-fertilization is an effective approach under salinity stress                                      | Sorour <i>et al.</i> (2022)   |
| Alfalfa ( <i>Medicago sativa</i> )  | Saline water (6 and 10 dS $\text{m}^{-1}$ )                 | Chemical and nano-fertilizers ( $\text{K}_2\text{SO}_4$ , ZnO and $\text{SiO}_2$ ) | The incorporation of nano-fertilizers with chemical fertilizers provide a promising strategy, especially in regions with low water quality                                                                                                                      | El-Shal <i>et al.</i> (2022)  |
| Soybean ( <i>Glycine max</i> )      | Salt affected soil (10.48 dS $\text{m}^{-1}$ )              | NPK nano-fertilizers, bio-fertilizers and humic acid                               | Integrated fertilization led to a slight decrease in soil pH, a significant decrease in EC values and significant increases in the availability of soil macro and micro-nutrients, as well as increases in the macro- and micronutrient contents in the grains. | Esmail <i>et al.</i> (2022)   |
| Barley ( <i>Hordeum vulgare</i> )   | Salt-affected soil and water                                | Phosphate solubilizing bacteria, nano phosphozink                                  | The integrated fertilization approach was effective in improving barley yield and enhancing soil physical and chemical properties.                                                                                                                              | Ibrahim and Hegab (2022)      |
| Wheat ( <i>Triticum aestivum</i> )  | Salt affected soil (6.33 to 8.13 dS $\text{m}^{-1}$ )       | Molasses, compost tea or K-humate combined with nano-silica                        | The co-application of organic substances and nano-silica was an effective approach to restore and conserve the soil and improve wheat productivity under salt-affected soil conditions                                                                          | Rashed <i>et al.</i> (2022)   |
| Cowpea ( <i>Vigna unguiculata</i> ) | Salt affected soil (8.85 dS $\text{m}^{-1}$ )               | Nano-NPK, organic fertilizers, NPK fertilization                                   | The combination of compost, chemical and nano-fertilizers improved soil nutrients' status and production of cowpea under saline soil conditions.                                                                                                                | Shaban <i>et al.</i> (2025)   |

#### 4. Crop production using integrated fertilizers in saline soils

##### 4.1. Benefits of Integrated Fertilization

Integrated fertilization offers numerous benefits and some limitations (**Figure 3**). This approach combines various agricultural practices, enhancing productivity and sustainability while addressing environmental concerns (Akter *et al.* 2025). The following points outline the key benefits and limitations of IFS:

- (1) Enhanced productivity: IFS promotes the synergistic integration of crops, livestock, and aquaculture, leading to increased overall productivity per unit area (Singh *et al.*, 2024; Bahadur *et al.*, 2024). It is worth to mention that, the integration of livestock and aquaculture, for example, is not a requirement of IFS. It may encourage farmers to be diverse in their production, but IFS can be done with only crops.
- (2) Soil health management: The balanced use of bio-, chemical, and nano-fertilizers improves soil fertility and maintains soil organic matter, crucial for sustainable crop production (Goda 2019; Hanurawaty, 2022).
- (3) Diversified income: By integrating multiple agricultural enterprises, farmers can generate diversified income streams throughout the year, reducing economic vulnerability (Singh *et al.*, 2024).
- (4) Environmental sustainability: Integrated practices reduce reliance on chemical fertilizers, thereby minimizing environmental degradation and greenhouse gas emissions (Hanurawaty, 2022).



**Fig. 3. An overview of integrated fertilization including benefits and limitations.**

##### 4.2. Limitations of Integrated Fertilization

Despite its limitations, integrated fertilization remains a promising strategy for enhancing agricultural sustainability and productivity. However, the complexity and resource demands may hinder its widespread adoption, particularly among smallholder farmers. The most important limits may include the following:

- (1) Complexity of management: Implementing IFS requires knowledge and skills to manage diverse components effectively, which can be challenging for smallholder farmers (Bahadur *et al.*, 2024).
- (2) Resource intensity: Initial investments in infrastructure and training can be high, potentially limiting adoption among resource-poor farmers (Hashmi and Saleem, 2017).
- (3) Dependence on external inputs: While integrated systems aim to reduce chemical use, they may still rely on external inputs, which can be economically burdensome (Goda, 2019).

##### 4.3. Integrated fertilizers and plant growth under saline soil conditions

The synergistic effects of combining bio-, mineral, organic, chemical, and nano- fertilizers on plant growth and nutrient uptake in saline soils have been extensively studied, revealing significant benefits for agricultural productivity. The integration of these fertilizers enhances soil properties, nutrient availability, and plant resilience against salinity, leading to improved crop yields, as follows:

### **(1) Enhanced Nutrient Availability**

The use of salt-tolerant plant growth-promoting bacteria (PGPB) like *Bacillus* species increases the availability of essential nutrients such as nitrogen, phosphorus, and potassium in saline soils (Mohaseb et al., 2019). Nano-fertilizers can improve nutrient solubility and uptake efficiency, allowing for controlled release and targeted delivery of nutrients, which is particularly beneficial in saline conditions (Tomar et al., 2024). At the same time, applying manures and composts as fertilizers improves the soil ecological environment through enhancing the nutrient availability (Tuo et al. 2025). Several reports have confirmed the role of nano-fertilizers in promoting soil nutrient availability (Rashid et al. 2023; Ansari et al. 2024).

### **(2) Improved Plant Growth**

Studies show that the combination of biofertilizers with chemical fertilizers, nanoparticles, and/or organic fertilizers significantly enhances plant growth parameters, including biomass yield and nutrient content in crops like rice and wheat (Mohaseb et al., 2019; Saleh et al., 2024). The interaction between nanoparticles and biofertilizers has been shown to promote better growth traits, such as increased chlorophyll content and reduced sodium accumulation in plant tissues (Karunakaran et al., 2024; Saleh et al., 2024). Different formulations of nano biofertilizers have been shown to be effective in enhancing plant growth (El-Bialy et al., 2023) and as bio-emerging strategies for sustainable agriculture (Garg et al. 2023; Sharma et al. 2023; Thirumurugan et al. 2024; Kumar et al. 2025).

### **(3) Soil Health and Salinity Mitigation**

The application of combined fertilizers leads to improved soil pH and reduced salinity levels, enhancing overall soil health and fertility (Saleh et al., 2024; Xing et al. 2025). Organic amendments, when combined with bio- and nano-fertilizers, promote salt leaching, further mitigating the adverse effects of salinity on plant growth (El-Sharkawy et al. 2024). While the synergistic effects of these fertilizers are promising, it is essential to consider the potential for over-reliance on such technologies, which may lead to unforeseen ecological impacts or nutrient imbalances if not managed properly.

### **(4) Mechanisms of synergistic interaction between nano-, bio-, organic and chemical fertilizers**

The synergistic interaction between nano, bio-, organic, and chemical fertilizers is facilitated by several underlying mechanisms that enhance nutrient efficiency and promote sustainable agricultural practices (**Figure 4**). This integration leverages the unique properties of nanoparticles, bioactive compounds, and traditional fertilizers to optimize plant growth and minimize environmental impact. This mechanism may include targeted delivery and controlled release of nutrients, enhanced nutrient bioavailability to cultivated plants, microbial synergy and physiological benefits (Yadav 2025). Nanofertilizers (NFs) enable precise nutrient delivery to plants, reducing nutrient losses and enhancing uptake efficiency through controlled release mechanisms (Fazelian and Morteza 2022; Padhan et al., 2023). The small size and high surface area of nanoparticles improve the solubility and bioavailability of nutrients, allowing for better absorption by plant roots (Padhan et al., 2023; Garg et al., 2023). Biofertilizers, when combined with NFs, promote beneficial microbial activity in the soil, enhancing nutrient cycling and plant health through mechanisms like rhizoremediation and disease resistance (Fazelian and Morteza 2022; Garg et al., 2023). The application of nano-biofertilizers can improve plant stress resistance and growth parameters, such as photosynthetic efficiency and leaf area, leading to increased crop yields (Kalenska et al., 2023; Rasool et al., 2024). While the integration of these fertilizers shows great promise for enhancing agricultural productivity, concerns regarding the long-term environmental impacts and potential toxicity of nanoparticles necessitate careful evaluation and regulation to ensure sustainable practices. The strong relationship between mineral and organic fertilizers for enriching the soil with fast and proper nutrients was considered by researchers several years ago. Thus, it is reported that the application of organo-mineral fertilizers can improve soil health and builds up a favorable environment for better plant growth, guarantee the slow release of nutrients, reduce nutrient leaching loss and greenhouse gas emissions than mineral fertilizers (Kafil Uddin et al. 2025). Organo-mineral fertilizers are also able to enhance crop productivity by increasing the nutrient use efficiency over mineral fertilizer application (Oliveira et al. 2025).




Fig. 4. The integration of applied fertilizers and environmental concern.

#### 4.4. Comparison of fertilizers

The use of fertilizers in agriculture is essential for maintaining soil fertility and ensuring crop productivity. However, the environmental and health impacts of different types of fertilizers, including organic, nano, bio-, and chemical fertilizers, have been a subject of extensive research (**Figure 5**). This section provides a comparison of the environmental and health risks associated with these fertilizers, as well as the regulatory frameworks governing their use (**Table 3**).

##### 4.4.1. Environmental risks of fertilizers

**(1) Chemical fertilizers:** Chemical fertilizers consist of synthetically formulated nutrient mixes and/or the transformation of mineral nutrient sources into forms that are more readily available to plants in the soil environment. Chemical fertilizers have been widely used in agriculture, particularly since World War II (Havlin et al., 2013). However, their production and use have significant environmental and human health impacts. The production of chemical fertilizers is energy-intensive and contributes to greenhouse gas emissions (Santolin et al., 2024). Additionally, the excessive use of nitrogen-based fertilizers has been linked to environmental pollution, including nitrate contamination of water bodies, a human and environmental health hazard (Brevik, 2010), and the release of nitrous oxide, a potent greenhouse gas (Huzenko, 2024). The leaching of nutrients from traditional fertilizers into water bodies can lead to eutrophication, which can have devastating effects on aquatic ecosystems (Eldridge et al., 2024; Jithendar et al., 2024). Furthermore, the use of traditional fertilizers has been associated with soil degradation, including acidification and the depletion of organic matter (Upadhyay et al., 2023).




Fig. 5. Comparative risk Analysis of Fertilizer types.

Table 3. A comparison among different types of fertilizers.

| Fertilizer Type      | Environmental Impact                                                                             | Health Impact                                                                                  | Regulatory Framework                                                                                |
|----------------------|--------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|
| Chemical fertilizers | High risk of nutrient leaching, soil degradation, and greenhouse gas emissions [1,2]             | Health risks from water and air pollution, and accumulation of toxic substances in food [2,3]  | Established regulatory frameworks, but need for stricter controls [1,2]                             |
| Nano-fertilizers     | Lower risk of nutrient leaching and runoff, but potential toxicity of nanoparticles [4,5]        | Potential health risks from nanoparticle toxicity [5,6]                                        | Regulatory frameworks in early stages of development, need for standardized testing protocols [5,6] |
| Bio-fertilizers      | Lower environmental impact, improves soil health, but potential for pathogen contamination [7,8] | Generally safer for human health, but risks from untreated waste and antibiotic resistance [9] | Established regulatory frameworks, but need for further research and harmonization [8,9]            |
| Organic fertilizers  | High environmental impact from mining and soil degradation [1,3]                                 | Health risks from toxic substances in soil and water [3]                                       | Regulatory frameworks in place, but need for stricter controls on mining and use [1,3]              |

List of refs. [1] Santolin et al. (2024), [2] Huzenko (2024), [3] Upadhyay et al. (2023), [4] Jithendar et al. (2024), [5] Noor & Elgharbawy (2024), [6] Langangmeilu et al. (2024), [7] Hu et al. (2024), [8] SAD (2024), [9] Gupta et al. (2024)

**(2) Nano fertilizers:** Nano fertilizers, which are fertilizers coated or encapsulated in nanomaterials, offer several environmental benefits compared to traditional fertilizers. These fertilizers release nutrients slowly, reducing the risk of leaching and runoff, which can contribute to water pollution (Jithendar et al., 2024). The controlled release of nutrients also reduces the need for frequent applications, thereby decreasing the overall environmental impact (Singh et al., 2023). However, the use of nano fertilizers is not without risks. The potential toxicity of nanoparticles to plants and soil organisms has been a concern. Some studies have shown that nanoparticles can cause phytotoxicity if applied in excess (Jithendar et al., 2024). Additionally, the long-term environmental fate of nanoparticles in agroecosystems is not yet fully understood, raising concerns about their potential accumulation in the environment (Noor and Elgharbawy, 2024).

**(3) Biofertilizers:** Biofertilizers focus on enhancement of the soil microbiome to enhance nutrient uptake and plant growth. They are considered more environmentally friendly than traditional fertilizers. These fertilizers improve soil health by increasing the availability of nutrients and enhancing soil organic matter (Hu et al., 2024). Biofertilizers also reduce the need for synthetic fertilizers, thereby decreasing greenhouse gas emissions and minimizing the risk of nutrient leaching (SAD 2024). However, the use of bio fertilizers is not without challenges. The production of bio fertilizers can be resource-intensive, and their effectiveness can vary depending on the specific microbial strains used (SAD 2024). Additionally, the use of bio fertilizers may require additional management practices, such as proper storage and application, to ensure their effectiveness and safety (Gupta et al., 2024).

**(4) Organic fertilizers:** Organic fertilizers are derived from organic sources such as animal waste and plant residues. They differ from biofertilizers in that the focus of organic fertilizers is on the addition of organic material to the soil with the benefit of slow nutrient release over time as the organic material decomposes, along with building of soil aggregates and its associated benefits (Liu et al. 2024; Tuo et al. 2025). While organic fertilizers also enhance soil microbial activity, unlike biofertilizers, organic fertilization does not involve adding particular target microorganisms to the soil (**Table 4**). The benefits of organic fertilizers include mitigation of greenhouse gas (GHG) emissions (He et al. 2023), improving soil health (Howe et al. 2024), sustainable remediation (Li et al. 2025), and novel fertilizers for organic farming (Balkrishna et al. 2025). Recently, a new approach towards application of organo-mineral fertilizer for improving soil health and building up a favorable environment for better plant growth was established (Kafil Uddin et al. 2025; Oliveira et al. 2025).

**Table 4. The fundamental differences between biofertilizers and organic fertilizers.**

| The comparison item               | Organic fertilizers                                                                                                                                 | Biofertilizers                                                                                                                                                                 |
|-----------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Definition                        | Decayed organic matter from plants and/or animals that directly enriches soil.                                                                      | Live microbes that enhance nutrient availability in soil.                                                                                                                      |
| Composition                       | Derived from decayed plant and/ or animal matter.                                                                                                   | Living microbes of bacteria, fungi, algae, solubilizing phosphorus or potassium                                                                                                |
| Examples                          | Farmyard manure, green manure, neem cake, poultry litter (compost, manure, bone meal, vermicompost, etc.).                                          | Rhizobium, Azotobacter, Azospirillum, Mycorrhizae, and phosphate-solubilizing bacteria (PSB).                                                                                  |
| Mode of action                    | Enhancing SOM, improving water retention, soil structure, microbial diversity, releasing nutrients slowly through microbial breakdown.              | By biological N-fixation, phosphate solubilization, or stimulating plant growth hormones (like IAA, cytokinins), and enhancing nutrient uptake efficiency.                     |
| Nutrient supply                   | Directly supply macronutrients (N, P, K) and micronutrients as they decompose, and acting as a slow-release fertilizer.                             | Enhancing nutrient availability (e.g., N-fixing bacteria convert atmospheric N <sub>2</sub> into plant-usable forms). More targeted in function (e.g., Rhizobium for legumes). |
| Application method and efficiency | Applied to soil by broadcasting, mixing, decomposition through microbes, provide SOM in slower nutrient release.                                    | Applied via seed treatment, root dipping or soil application, for microbial survival specific conditions (i.e., moisture, pH, host compatibility) are required.                |
| Environmental dimension           | Improving soil health but may introduce pathogens if not properly composted, excessive use can lead to nutrient leaching (e.g., nitrate pollution). | Eco-friendly, reduce dependency on synthetic fertilizers, do not cause pollution but may have limited shelf life.                                                              |
| Shelf life and storage            | Longer shelf life if stored properly (no moisture).                                                                                                 | Short shelf life (6–12 months) due to living microbes; require proper storage (cool, dry conditions).                                                                          |

Sources: Lal (2025), El-Ghamry et al. (2018), Lio et al. (2025)

#### 4.4.2. Health risks of fertilizers

**(1) Chemical fertilizers:** The excessive use of traditional chemical fertilizers has been linked to several health risks (Steffan et al., 2018). The contamination of water bodies with nitrates from nitrogen-based fertilizers can lead to health problems, including methemoglobinemia in infants (Huzenko, 2024). Additionally, the release of ammonia from fertilizers can contribute to air pollution, which has been linked to respiratory and cardiovascular diseases (Huzenko, 2024). The use of traditional chemical fertilizers has also been associated with the accumulation of toxic substances in food crops, which can pose health risks to consumers (Upadhyay et al., 2023). Furthermore, the use of chemical fertilizers has been linked to the development of antibiotic resistance in soil microorganisms, which can have broader implications for human health (Gupta et al., 2024). Chemical fertilizers are generally considered safe for human health when used properly. However, the excessive use of these fertilizers can lead to the accumulation of toxic substances in soil and water, which can pose health risks to humans, animals and the broader environment (Upadhyay et al., 2023). Additionally, the mining to obtain raw materials for the manufacture of chemical fertilizers can lead to the release of heavy metals and other toxic substances into the environment, which can have long-term effects on human health (Steffan et al., 2018; Santolin et al., 2024).

**(2) Nanofertilizers:** The health risks associated with nano fertilizers are primarily related to the potential toxicity of nanoparticles. Some studies have shown that nanoparticles can cause oxidative stress and DNA damage in plants and soil organisms (Noor & Elgharbawy, 2024). Additionally, the inhalation of nanoparticles by humans can lead to respiratory and other health problems (Langangmeilu et al., 2024). However, the health risks associated with nano fertilizers are still not fully understood, and more research is needed to assess their long-term effects on human health and the environment (Noor and Elgharbawy, 2024). Regulatory frameworks governing the use of nano fertilizers are still in the early stages of development, and there is a need for standardized testing protocols to ensure their safe use (Langangmeilu et al., 2024).

**(3) Bio-fertilizers:** Bio-fertilizers are generally considered safer for human health and the environment compared to chemical fertilizers. These fertilizers are derived from natural sources and do not contain synthetic chemicals, which reduces the risk of toxic substances accumulating in food crops (SAD 2024). Additionally, biofertilizers can improve soil health, which can lead to the production of healthier food crops (Hu et al., 2024). However, the use of bio fertilizers can also pose some health risks if not managed properly. For example, the use of untreated animal waste as a bio fertilizer can lead to the contamination of soil and water with pathogens, which can pose health risks to humans and animals (Gupta et al., 2024). Additionally, the use of certain microbial inoculants in bio fertilizers can lead to the development of antibiotic resistance in soil microorganisms, which can have broader implications for human health (Gupta et al., 2024).

**(4) Organic fertilizers:** the use of untreated animal waste as an organic fertilizer can lead to the contamination of soil and water with pathogens, which can pose health risks to humans and animals (Gupta et al., 2024). Additionally, the use of certain microbial inoculants in bio fertilizers can lead to the development of antibiotic resistance in soil microorganisms, which can have broader implications for human health (Gupta et al., 2024).

#### 4.4.3. Regulatory frameworks governing fertilizer use

**(1) Chemical fertilizers:** The use of traditional fertilizers is governed by various regulatory frameworks aimed at minimizing their environmental and health impacts. For example, the European Union has implemented regulations to reduce the use of nitrogen-based fertilizers and to promote the use of organic fertilizers (Santolin et al., 2024). Additionally, many countries have established standards for the safe use of fertilizers, including guidelines for application rates and the management of fertilizer waste (Huzenko, 2024).

**(2) Nanofertilizers:** The regulatory frameworks governing the use of nano fertilizers are still in the early stages of development. There is a need for standardized testing protocols to assess the safety and efficacy of nano fertilizers, as well as their potential environmental and health impacts (Noor & Elgharbawy, 2024). Some countries have established regulatory frameworks for the use of nanotechnology in agriculture, but these frameworks are not yet harmonized at the international level (Langangmeilu et al., 2024).

**(3) Bio- and organic fertilizers:** The use of bio-and organic fertilizers is governed by regulatory frameworks aimed at ensuring their safety and efficacy. For example, the European Union has established regulations for the use of microbial inoculants in biofertilizers, including requirements for the safety of microorganisms and their potential impact on the environment (SAD 2024).

Additionally, many countries have established standards for the production and use of bio fertilizers, including guidelines for quality control and labeling (Gupta *et al.*, 2024). Regulation 2019/1009 is well-known and issued by the European Union (EU)-Fertilising Products Regulation, which aimed to create a single market for organic fertilizer products, ensuring safety, quality, and appropriate labeling while also addressing environmental and health concerns (EU 2022). This regulation included some rules to open a single market to organic and waste-based fertilizers, to establish limit values for toxic pollutants in fertilizing products, and to allow optional harmonization (EU 2022). A very wide range of fertilizing products within the EU including organo-mineral fertilizers, organic fertilizers, and biostimulants is reported (EU 2025).

## 5. Conclusions

As a result of climate change, soil salinization may be accelerated with increased pressure on global food production. This needs continuous innovative solutions to mitigate and treatment of such global problems. Soil salinity can be found in new places all over the world due to this change. IFS is an important approach for mitigating soil salinity, but to what extent can this agronomic strategy be followed? More studies are needed to emphasize the right combinations of different kinds of fertilizers in various agroecological systems. The most important concern is to reduce the use of chemical fertilizers by replacing them with organic, nano, and biofertilizers when this is appropriate and effective, and in finding ways to maximize the contributions of all types of fertilizers through the use of IFS.

## Declarations

### Ethics approval and consent to participate

**Author Contributions:** All authors helped prepare the MS and agree to publish it.

**Funding:** There is no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** There is no conflict of interest among the authors.

## 6. References

- Absamat o'g'li, E. U., Zhou, J. (2022). Bio-Organic Fertilizer Combined with Different Amendments Improves Nutrient Enhancement and Salt Leaching in Saline Soil: A Soil Column Experiment. *Water*, 14(24), 4084. <https://doi.org/10.3390/w14244084>
- Akter S, Ahmed JU, Begum IA, Sarkar MA, Fatema K, Mahmud A, Ding S, Alam MJ (2025). Integrated farming system-A means of improving farmers' well-being in the wetland ecosystem of Bangladesh. *Farming System*, 3, Issue 1, 100127. <https://doi.org/10.1016/j.farsys.2024.100127>.
- Ali, M.Y.; Sina, A.A.I.; Khandker, S.S.; Neesa, L.; Tanvir, E.M.; Kabir, A.; Khalil, M.I.; Gan, S.H (2021). Nutritional Composition and Bioactive Compounds in Tomatoes and Their Impact on Human Health and Disease: A Review. *Foods* 10, 45. <https://dx.doi.org/10.3390/foods10010045>
- Alotaibi, M. M., Aljuaid, A., Alsudays, I. M., Aloufi, A. S., AlBalawi, A. N., Alasmari, A., et al. (2024). Effect of bio-fertilizer application on agronomic traits, yield, and nutrient uptake of Barley (*Hordeum vulgare*) in saline soil. *Plants*, 13(7), 951.
- Al-Taey, D. K., Al-Musawi, Z. J. (2019). Effect of Nano-fertilizers, salicylic acid, and organic matter in growth and yield of rocket (*Eruca sativa* Mill) under Salt stress. *International Journal of Botany Studies*, 4(3), 77-81.
- Ansari MM, Shin M, Kim M, Ghosh M, Kim SH, Son YO (2024). Nano-enabled strategies in sustainable agriculture for enhanced crop productivity: A comprehensive review. *J Environ Manage.*, 372:123420. doi: 10.1016/j.jenvman.2024.123420.
- Ashford, E. H. (2023). *Management of Salt-Affected Soils for Increasing Crop Productivity* (pp. 113–121). [https://doi.org/10.1007/978-981-19-9159-2\\_7](https://doi.org/10.1007/978-981-19-9159-2_7)
- Awwad, E. A., Mohamed, I. R., Abd El-Hameed, A. M., Zaghloul, E. A. (2022). The Co-addition of soil organic amendments and natural bio-stimulants improves the production and defenses of the wheat plant grown under the dual stress of salinity and alkalinity. *Egyptian Journal of Soil Science*, 62(2), 137-153.

- Bahadur, R., Kumar, N., Tripathi, U., Singh, A. K., Kumari, G., Alagesan, A., Senthilkumar, T., Malathi, G., Prabhavathi, S. J. (2024). Enhancing Productivity and Sustainability through Integrated Farming Systems: A Review. *International Journal of Environment and Climate Change*. <https://doi.org/10.9734/ijecc/2024/v14i13887>
- Baig, K., Ali, Z., Warraich, I., Kausar, R., Kashaf, A., Gul, N., Khan, O., Nawaz, R., Ahmad, H. (2023). *Sustainable cotton cultivation in saline soils: challenges, innovations, and future prospects*. <https://doi.org/10.54112/bcsrj.v2023i1.584>
- Balkrishna A, Kaushik P, Singh S, Agrahari P, Kumar B, Kumar P, Arya VP (2025). Potential use of sewage sludge as fertilizer in organic farming. *Cleaner Waste Systems*, Volume 10, 100245. <https://doi.org/10.1016/j.clwas.2025.100245>.
- Chen YN, Fan C, Šereš M, Šerešová M, Vymazal J, Pan SY (2025). Lifecycle environmental benefits of integrated rational fertilization, biochar, and constructed wetland in mitigating nutrient loading. *Agricultural Water Management*, 307, 109202. <https://doi.org/10.1016/j.agwat.2024.109202>.
- Chi Y, Fan M, Zhang Z, Qu Y (2025). Zoning the soil salinization levels in the northern China's coastal areas based on high-resolution soil mapping. *Ecological Indicators*, 172, 113303. <https://doi.org/10.1016/j.ecolind.2025.113303>.
- Choudhary, O. P., Bajwa, M. S., Josan, A. S. (2003). Fertilizer Management in Salt-Affected Soils: A Review. *Journal of Research*, 40(2), 153–171. <http://www.indianjournals.com/ijor.aspx?target=ijor:jre&volume=40&issue=2&article=004&type=pdf>
- Dai S, Feng, W, Song, F, Li, T, Tao, Y, Yang, F, Miao, Q, Duan, P, Liao, H, Shi, H, Gonçalves, JM, Duarte IM (2025). Review of biological algal fertilizer technology: Alleviating salinization, sequestering carbon, and improving crop productivity, *Bioresource Technology*, 429, 132507. <https://doi.org/10.1016/j.biortech.2025.132507>.
- Duan H, Gao R, Liu X, Zhang L, Wang Y, Jia X, Wang X, Zheng S, Jing Y (2025). The coupling of straw, manure and chemical fertilizer improved soil salinity management and microbial communities for saline farmland in Hetao Irrigation District, China. *Journal of Environmental Management*, 380, 124917. <https://doi.org/10.1016/j.jenvman.2025.124917>.
- El-Ghamry, A., Mosa, A., Alshaal, T., El-Ramady, H. (2018). Nanofertilizers vs. Biofertilizers: New Insights. *Environment, Biodiversity and Soil Security*, 2(2018), 51-72. doi: 10.21608/jenvbs.2018.3880.1029
- El-Ramady, H., Faizy, S., Amer, M., Elsakhawy, T., Omara, A. E., Eid, Y., Brevik, E. (2022). Management of Salt-Affected Soils: A Photographic Mini-Review. *Environment, Biodiversity and Soil Security*, 6(2022), 61-79. doi: 10.21608/jenvbs.2022.131286.1172
- El-Ramady, H., Prokisch, J., Mansour, H., Bayoumi, Y. A., Shalaby, T. A., Veres, S., Brevik, E. C. (2024). Review of Crop Response to Soil Salinity Stress: Possible Approaches from Leaching to Nano-Management. *Soil Systems*, 8(1), 11. <https://doi.org/10.3390/soilsystems8010011>
- El-Shal, R. M., El-Naggar, A. H., El-Beshbesy, T. R., Mahmoud, E. K., El-Kader, N. I. A., Missaui, A. M., et al. (2022). Effect of nano-fertilizers on alfalfa plants grown under different salt stresses in hydroponic system. *Agriculture*, 12(8), 1113.
- Esmaeil, M. A., Shaban, K. A., Fattah, A. K. A., Mohammed, I. A. B. E. I. (2022). Assessment of the Efficiency of Nano, Bio and Organic Fertilizers on Soil Chemical Properties and Soybean Productivity in Saline Soil. *International Journal of Plant and Soil Science*, 34(17), 24-38.
- EU (2022). New EU rules prepare the ground for more use of organic and waste-based fertilisers. [https://single-market-economy.ec.europa.eu/news/new-eu-rules-prepare-ground-more-use-organic-and-waste-based-fertilisers-2022-07-15\\_en](https://single-market-economy.ec.europa.eu/news/new-eu-rules-prepare-ground-more-use-organic-and-waste-based-fertilisers-2022-07-15_en). Accessed on July 15, 2025
- EU (2025). EU Fertilizing Product Regulation. <https://www.fertilizerseurope.com/agriculture-environment/fertilizing-products-regulation/> Accessed on July 15, 2025
- Faisal, M., Raza, A., Batool, A., Zeshan, M., Ismail, M., Farman, M., Anwar, I., Mobeen, A., Dilshad, M. (2024). Unveiling the Impacts of Soil Salinity: Challenges, Mechanisms, and Sustainable Solutions. *Scholars Journal of Agriculture and Veterinary Sciences*, 11(08), 152–162. <https://doi.org/10.36347/sjavs.2024.v11i08.002>
- Farias GD, Bremm C, Lemaire G, Assmann TS, Simões VJLP, Tales Tiecher, Alves LA, Martins AP, Carvalho PF (2025). Can system fertilization improve plant nutrition and ensure nutrient carryover to successive crops in integrated crop-livestock systems? *Field Crops Research*, 109985. <https://doi.org/10.1016/j.fcr.2025.109985>.

- Fazelian N, Y Morteza (2022). Nano-biofertilizers for enhanced nutrient use efficiency. In: Nano-enabled Agrochemicals in Agriculture, Editor(s): Mansour Ghorbanpour, Muhammad Adnan Shahid, Academic Press, Pp: 145-158, Elsevier eBooks. <https://doi.org/10.1016/b978-0-323-91009-5.00023-9>
- Garg, D., Sridhar, K., Inbaraj, B. S., Chawla, P., Tripathi, M., Sharma, M. (2023). Nano-Biofertilizer Formulations for Agriculture: A Systematic Review on Recent Advances and Prospective Applications. *Bioengineering*, 10. <https://doi.org/10.3390/bioengineering10091010>
- Goda, D. D. (2019). Effect of Integrated Soil Fertility Management on Crop Production: A Review. *Journal of Biology, Agriculture and Healthcare*, 9(8), 1–8. <https://iiste.org/Journals/index.php/JBAH/article/download/47335/48879>
- Gupta, A., Saharan, B. S., Sadh, P. K., Duhan, J. S. (2024). Green to green: harnessing bio-fertilizers for sustainable pulse production and food security. *Discover Plants.*, 1(1). <https://doi.org/10.1007/s44372-024-00079-y>
- Hanurawaty, N. Y. (2022). Integrated Soil Fertility Management for Climate Change Mitigation and Agricultural Sustainability (pp. 129–154). CRC Press eBooks. <https://doi.org/10.1201/9781003175018-8>
- Hasan, B. K., Saad, T. M. (2020). Effect of Nano biological and mineral fertilizers on NPK uptake in wheat (*Triticum aestivum* L.). *Indian Journal of Ecology*, 47(12), 126-130.
- Hashmi, S., Saleem, Q. (2017). Best Practices for Sustaining Agriculture: Integrating Indigenous and Modern Methods of Soil Fertility Management Inbeed District. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 11(05), 49–52. <https://doi.org/10.9790/2402-1105044952>
- He Z, Ding B, Pei S, Cao H, Liang J, Li Z (2023). The impact of organic fertilizer replacement on greenhouse gas emissions and its influencing factors. *Sci Total Environ.* 905:166917. doi: 10.1016/j.scitotenv.2023.166917.
- Hou, P., Li, B., Cao, E., Jian, S., Liu, Z., Li, Y., et al. (2025). Optimizing maize yield and mitigating salinization in the Yellow River Delta through organic fertilizer substitution for chemical fertilizers. *Soil and Tillage Research*, 249, 106498.
- Howe JA, McDonald MD, Burke J, Robertson I, Coker H, Gentry TJ, Lewis KL (2024). Influence of fertilizer and manure inputs on soil health: A review. *Soil Security*, 16, 100155. <https://doi.org/10.1016/j.soisec.2024.100155>.
- Hu, M.-C., Xue, H., Wade, A. J., Gao, N., Qiu, Z., Long, Y., Shen, W.-S. (2024). Biofertilizer supplements allow nitrogen fertilizer reduction, maintain yields, and reduce nitrogen losses to air and water in China paddy fields. <https://doi.org/10.1016/j.agee.2023.108850>
- Huzenko, M. (2024). Interrelations of Current Health Expenditure and Fertilizer Consumption in European Countries: Step-by-Step to Sustainable Agricultural Practices. *Health Economics and Management Review*, 5(4), 1–18. <https://doi.org/10.61093/hem.2024.4-01>
- Huzenko, M. (2024). Interrelations of Current Health Expenditure and Fertilizer Consumption in European Countries: Step-by-Step to Sustainable Agricultural Practices. *Health Economics and Management Review*, 5(4), 1–18. <https://doi.org/10.61093/hem.2024.4-01>
- Ibrahim, G. A., Hegab, R. (2022). Improving yield of Barley using bio and nano fertilizers under saline conditions. *Egyptian Journal of Soil Science*, 62(1), 41-53.
- Jia F, Wenhao Li, Bo Zhou, Yang Xiao, Buchun Si, Aizimuhan Saikeshan, Tingbo Lv, Zhenhua Wang Z (2025). Consecutive high-efficient water-saving irrigation increase crop yield and decrease soil salinity through reconstructing rhizosphere soil bacterial communities. *Science of The Total Environment*, 959, 178238. <https://doi.org/10.1016/j.scitotenv.2024.178238>.
- Jithendar, B., Kumar, R., Rana, N. (2024). Revolutionizing Crop Nutrition: Exploring Nano Fertilizers in Agriculture. *International Journal of Plant & Soil Science* 36 (6):327-39. <https://doi.org/10.9734/ijpss/2024/v36i64635>
- Kafil Uddin M, Saha BK, Wong VNL, Patti AF (2025). Organo-mineral fertilizer to sustain soil health and crop yield for reducing environmental impact: A comprehensive review. *European Journal of Agronomy*, Volume 162, 127433. <https://doi.org/10.1016/j.eja.2024.127433>.
- Kalenska, S., Novytska, N., Kalenskyi, V., El Farouk Osman, O., & Melnychenko, V. (2023). Efficiency of the compensatory scheme of using Organic Acid Nano-Complex microfertilizers in spring wheat cultivation technology. *Рослинництво Та Ґрунтознавство*. <https://doi.org/10.31548/plant4.2023.61>

- Karunakaran, A., Fathima, Y., Singh, P., Beniwal, R., Singh, J., Ramakrishna, W. (2024). Next-Generation Biofertilizers: Nanoparticle-Coated Plant Growth-Promoting Bacteria Biofertilizers for Enhancing Nutrient Uptake and Wheat Growth. *Agriculture*, 14(4), 517. <https://doi.org/10.3390/agriculture14040517>
- Kaviyazhagan, S., Prakash, K. (2023). AGRONOMIC VIEW OF SOIL SALINITY, ITS EFFECTS AND RECLAMATION - A Review. *Madras Agricultural Journal*, 110, 1–5. <https://doi.org/10.29321/maj.10.20008>
- Lal, R. (2015). Restoring Soil Quality to Mitigate Soil Degradation. *Sustainability*, 7(5), 5875–5895. DOI:10.3390/su7055875
- Langangmeilu, G., Heisnam, P., Moirangthem, A., Vani, N. (2024). Nanotechnology in agriculture: opportunity and challenges (pp. 257–265). <https://doi.org/10.58532/v3becs24p5ch1>
- Li C, Cai H, Liu Y (2025). Sustainable approaches to tailings remediation: Role of organic fertilizers. *Environmental Research*, 276, 121454. <https://doi.org/10.1016/j.envres.2025.121454>.
- Li G, Shan Y, Nie W, Sun Y, Su L, Mu W, Qu Z, Yang T, Wang Q (2024). Application of carboxymethyl cellulose sodium (CMCNa) in maize-wheat cropping system (MWCS) in coastal saline-alkali soil. *Sci Total Environ*. 912:169214. doi: 10.1016/j.scitotenv.2023.169214.
- Li T, Wang S, Liu S, Zhang X, Dong H, Dai S, Chai L, Li H, Lv Y, Li T, Gao Q, Li G, Ma X (2025). Trade-offs of organic amendment input on soil quality and crop productivity in saline-alkali land globally: A meta-analysis. *European Journal of Agronomy*, 164, 127471. <https://doi.org/10.1016/j.eja.2024.127471>.
- Li Z, Meng Q, Li L, Bai Z, Li Y, Liu h, Li P, Wang T (2025). Integrated deep vertical rotary tillage and subsurface pipe drainage techniques for sustainable soil salinization management and cotton production in arid regions. *Agricultural Water Management*, 312, 109429. <https://doi.org/10.1016/j.agwat.2025.109429>.
- Liao Z, Boubakri H, Chen B, Farooq M, Lai Z, Kou H, Fan J (2025). Biofertilizers as an eco-friendly approach to combat drought stress in plants. *Biocatalysis and Agricultural Biotechnology*, Volume 64, 103510. <https://doi.org/10.1016/j.bcab.2025.103510>
- Liu M, Wang X, Yang B, Song D, Xu X, Ding W, He P, Zhou W (2025). Optimized fertilization strategies enhanced integrated benefits in nutrient management for open-field cabbage production. *Journal of Agriculture and Food Research*, 20, 101801. <https://doi.org/10.1016/j.jafr.2025.101801>.
- Liu x, Zhang Y, Wang Z, Chen Z (2024). The contribution of organic and chemical fertilizers on the pools and availability of phosphorus in agricultural soils based on a meta-analysis. *European Journal of Agronomy*, 156, 127144. <https://doi.org/10.1016/j.eja.2024.127144>.
- Liu Y, Ze Zhang, Xiangxiang Wang, Ruiqiao Wu, Shuai Ding, Shuang Wang, Jianping Chen, Tida Ge, Zhu Z (2025). Gypsum application increases microbial activity and organic carbon mineralization in saline paddy soils. *Applied Soil Ecology*, 208, 106004. <https://doi.org/10.1016/j.apsoil.2025.106004>.
- Liu Z, Gao C, Yan Z, Shao L, Chen S, Niu J, Zhang X (2024). Effects of long-term saline water irrigation on soil salinity and crop production of winter wheat-maize cropping system in the North China Plain: A case study. *Agricultural Water Management*, 303, 109060. <https://doi.org/10.1016/j.agwat.2024.109060>.
- Liu, J., Xie, W., Yang, J., Yao, R., Wang, X., Li, W. (2023). Effect of different fertilization measures on soil salinity and nutrients in salt-affected soils. *Water*, 15(18), 3274.
- Liu, Y., Lan, X., Hou, H., Ji, J., Liu, X., Lv, Z. (2024). Multifaceted Ability of Organic Fertilizers to Improve Crop Productivity and Abiotic Stress Tolerance: Review and Perspectives. *Agronomy*, 14(6), 1141. <https://doi.org/10.3390/agronomy14061141>
- Majeed, A., Muhammad, Z. (2019). Salinity: A Major Agricultural Problem—Causes, Impacts on Crop Productivity and Management Strategies (pp. 83–99). Springer, Cham. [https://doi.org/10.1007/978-3-030-06118-0\\_3](https://doi.org/10.1007/978-3-030-06118-0_3)
- Melebari, D. (2025). The use of soil amendments and foliar application can improve plant production under salinity stress conditions. *Egyptian Journal of Soil Science*, 65(1): 321–338.
- Mishra AK, Das R, George Kerry R, Biswal B, Sinha T, Sharma S, Arora P and Kumar M (2023) Promising management strategies to improve crop sustainability and to amend soil salinity. *Front. Environ. Sci.* 10:962581. doi: 10.3389/fenvs.2022.962581

- Mohaseb, M. I., Kenawy, M. H. M., Shaban, K. A. H. (2019). Role of Mineral and Bio-fertilizers on Some Soil Properties and Rice Productivity under Reclaimed Saline Soils. 1–12. <https://doi.org/10.9734/ASRJ/2019/V2I130041>
- Mukhopadhyay R, Sarkar B, Jat HS, Sharma PC, Bolan NS. Soil salinity under climate change: Challenges for sustainable agriculture and food security. *J Environ Manage*. 2021, 280:111736. doi: 10.1016/j.jenvman.2020.111736.
- Noor, N., Elgharrawy, A. A. M. (2024). Fate of nanofertilizer in agroecosystem (pp. 281–295). Elsevier BV. <https://doi.org/10.1016/b978-0-443-15570-3.00021-1>
- Oliveira RC, Alves MF, Luz JMQ, Blank AF, Nizio DAC, Nogueira PCL, Silva SM, Castoldi R (2025). Organo-Mineral Fertilizer Improves *Ocimum basilicum* Yield and Essential Oil. *Plants (Basel)*. 14(7):997. doi: 10.3390/plants14070997.
- Padhan, S. R., Kar, I., Mohanty, A., Panigrahi, K. K. (2023). Nanofertilizers: Types, Synthesis, Methods, and Mechanisms (pp. 61–98). Springer International Publishing. [https://doi.org/10.1007/978-3-031-41329-2\\_3](https://doi.org/10.1007/978-3-031-41329-2_3)
- Rashed, S. H., Sorour, S., Aelhag, D., Amer, M. M. (2022). Effect of some organic substances and foliar application of nano-silica on physico-chemical soil properties and yield of wheat in salt affected soils. *International Journal of Plant & Soil Science*, 5, 15-27.
- Rashid MI, Shah GA, Sadiq M, Amin NU, Ali AM, Ondrasek G, Shahzad K (2023). Nanobiochar and Copper Oxide Nanoparticles Mixture Synergistically Increases Soil Nutrient Availability and Improves Wheat Production. *Plants (Basel)*, 12(6):1312. doi: 10.3390/plants12061312.
- Rasool, A., Herwahu Krismastuti, F. S., Zulfajri, M., Meliana, Y., Sudewi, S. (2024). A smart way to increase the growth and productivity of crops through nano-fertilizer (pp. 333–346). Elsevier BV. <https://doi.org/10.1016/b978-0-323-95721-2.00013-0>
- Rolón-Cárdenas GA, Rodríguez-González V (2025). Physiological response of tomato plants grown in saline and non-saline soils to foliar application of SiO<sub>2</sub>/TiO<sub>2</sub> nanocomposites. *Biocatalysis and Agricultural Biotechnology*, 65, 103536. <https://doi.org/10.1016/j.bcab.2025.103536>.
- SAD, Sustainable Agriculture Development (2024). The Role of Biofertilizers in Soil Fertility and Crop Yield Improvement. 2(1), 1–5. <https://doi.org/10.25163/agriculture.2110005>
- Saleh, M. A., Metwally, S. M., Ayman, M., Abdelghany, A. M. (2024). Synergistic Approaches for Wheat Productivity in Saline-Sodic Soils: Nano-Gypsum, Bacillus Inoculation and Nanomaterial Foliar Treatments. *Deleted Journal*, 1(1), 36–55. <https://doi.org/10.21608/bct.2024.399466>
- Santolin, J., Vlaeminck, S. E., Appiah-Twum, H., Van Winckel, T., Spiller, M. (2024). Consequential LCA between NPK fertilizers from microbial, animal, plant, and mineral origin highlights resource constraints and environmental impacts. *Journal of Cleaner Production*. <https://doi.org/10.1016/j.jclepro.2024.142312>
- Santolin, J., Vlaeminck, S. E., Appiah-Twum, H., Van Winckel, T., Spiller, M. (2024). Consequential LCA between NPK fertilizers from microbial, animal, plant, and mineral origin highlights resource constraints and environmental impacts. *Journal of Cleaner Production* <https://doi.org/10.1016/j.jclepro.2024.142312>
- Shaban, K. A., I Gaber, E. S., Soliman, A. S., Ozraeil, H M. (2025). Impact of A Nano-NPK Fertilizer With/Without The Compost on Some Soil Chemical Characteristics and Cowpea Yield under Salinity Conditions. *Egyptian Journal of Soil Science*, 65(2).
- Sheoran, H. S. (2023). Fertility Management in Salt Affected Soils under Rice-wheat Cropping System: A Review. *Agricultural Reviews*. <https://doi.org/10.18805/ag.r-2615>
- Shoukat, A., Maryam, U., Pitann, B., Zafar, M. M., Nawaz, A., Hassan, W., Seleiman, M. F., Saqib, Z. A., Mühling, K. H. (2025). Efficacy of Nano and Conventional Zinc and Silicon Fertilizers for Nutrient Use Efficiency and Yield Benefits in Maize Under Saline Field Conditions. *Plants*, 14(5), 673. <https://doi.org/10.3390/plants14050673>
- Shoukat, A., Saqib, Z. A., Akhtar, J., Aslam, Z., Pitann, B., Hossain, M. S., Mühling, K. H. (2024). Zinc and silicon nano-fertilizers influence ionic and metabolite profiles in maize to overcome salt stress. *Plants*, 13(9), 1224.
- Simões VJLP, Duarte LP, Rintzel RDD, Martins AP, Tiecher T, Mori LD, Bremm C, Carneiro MAC, Carvalho PCF (2025). System fertilization improves soil quality and increases primary production in an integrated crop-livestock system. *Journal of Integrative Agriculture*. <https://doi.org/10.1016/j.jia.2025.03.002>.

- Singh, A., Rajput, V. D., Tawaha, A. M., Al Zoubi, O. M., Habeeb, T. H., Rawat, S., Ghazaryan, K., Minkina, T. (2023). A Review on Crop Responses to Nanofertilizers for Mitigation of Multiple Environmental Stresses. <https://doi.org/10.12912/27197050/169313>
- Singh, J., Singh, B., Singh, A. K., Rani, P. (2024). Integrated Farming System: A Sustainable Approach towards Modern Agriculture. *International Journal of Plant and Soil Science*, 36(9), 641–649. <https://doi.org/10.9734/ijpss/2024/v36i95012>
- Song Y, Sun J, Cai M, Li J, Bi M, Gao M (2025). Effects of management of plastic and straw mulching management on crop yield and soil salinity in saline-alkaline soils of China: A meta-analysis. *Agricultural Water Management*, 308, 109309. <https://doi.org/10.1016/j.agwat.2025.109309>.
- Sorour, S., Amer, M. M., El Hag, D., Hasan, E. A., Awad, M., Kizilgeci, F., et al. (2021). Organic amendments and nano-micronutrients restore soil physico-chemical properties and boost wheat yield under saline environment. *Fresenius Environmental Bulletin*, 30(9), 10941-10950.
- Su X, Zhang L, Meng H, Zhao J, Wang H, Shang L, Guo Q, Xing Y, Song X, Sun X, Zhang X, Tian X, Mao L (2025). The different pathways of microbial regulation of organic carbon turnover in the topsoil and subsoil of coastal saline soil after long-term stubble return and subsoiling. *Applied Soil Ecology*, 208, 105970. <https://doi.org/10.1016/j.apsoil.2025.105970>.
- Tarolli P, Luo J, Park E, Barcaccia G, Masin R (2024). Soil salinization in agriculture: Mitigation and adaptation strategies combining nature-based solutions and bioengineering. *iScience.*, 27(2):108830. doi: 10.1016/j.isci.2024.108830.
- Tomar, B. S., Patle, T., Parihar, S. S., Singh, P., Tomar, S. S. (2024). Enhancing Nutrient Uptake With Nano Fertilizers and Soil Amendments (pp. 43–59). IGI Global. <https://doi.org/10.4018/979-8-3693-1890-4.ch003>
- Tuo Y, Xie C, Wang S, Xiang P, Yang Q, He X (2025). Irrigation combined with organic fertilizer could promote the environmental health of soil available nutrients and improve the functional activity of saponins in *Panax Notoginseng*. *Agricultural Water Management*, 313, 109444. <https://doi.org/10.1016/j.agwat.2025.109444>.
- Upadhyay, P. K., Dey, A., Singh, V. K., Dwivedi, B. S., Babu, V. R., Rathore, S. S., Singh, R. K., Shekhawat, K., Yadav, D., Singh, D., Shukla, G. (2023). Conjoint application of nano-urea with conventional fertilizers: An energy efficient and environmentally robust approach for sustainable crop production. *PLOS ONE*, 18(7), e0284009. <https://doi.org/10.1371/journal.pone.0284009>
- Upadhyay, P. K., Dey, A., Singh, V. K., Dwivedi, B. S., Babu, V. R., Rathore, S. S., Singh, R. K., Shekhawat, K., Yadav, D., Singh, D., Shukla, G. (2023). Conjoint application of nano-urea with conventional fertilizers: An energy efficient and environmentally robust approach for sustainable crop production. *PLOS ONE*, 18(7), e0284009. <https://doi.org/10.1371/journal.pone.0284009>
- Wang T, Jiawei Jing, Pengfei Huang, Xinyu Guo, Chuan Li, Qu Y (2025). Bioremediation of alkane-containing saline soils using the long-chain alkane-degrading bacterium *Pseudomonas aeruginosa* DL: Effects, communities, and networks. *Journal of Hazardous Materials*, 488, 137401. <https://doi.org/10.1016/j.jhazmat.2025.137401>.
- Wang Y, Guo T, Tian C, Zhang K, Zhao Z, Mao X, Mai W (2025). Effects of root growth on salt leaching and soil structure improvement in saline soils: A case study of *Suaeda salsa*. *Agricultural Water Management*, 314, 109533. <https://doi.org/10.1016/j.agwat.2025.109533>.
- Wang, M., Wang, R., Sun, Q., Li, Y., Xu, L., Wang, Y. (2024). Integrated Drip Irrigation Regulates Soil Water–Salt Movement to Improve Water Use Efficiency and Maize Yield in Saline–Alkali Soil. *Water*, 16(17), 2509. <https://doi.org/10.3390/w16172509>
- Wang, N., Zhang, T., Li, Y., Cong, A., Lian, J., Feng, K. (2025). Integrated application of fertilization increased maize (*Zea mays* L.) yield by improving soil quality, particularly under limited water conditions in a semi-arid sandy area. *Agricultural Water Management*, 309, 109334.
- Wei, X., Xie, B., Wan, C., Song, R., Zhong, W., Xin, S., Song, K. (2024). Enhancing Soil Health and Plant Growth through Microbial Fertilizers: Mechanisms, Benefits, and Sustainable Agricultural Practices. *Agronomy*, 14(3), 609. <https://doi.org/10.3390/agronomy14030609>
- Wu Q, Chen Y, Dou X, Liao D, Li K, An C, Li G, Dong Z. Microbial fertilizers improve soil quality and crop yield in coastal saline soils by regulating soil bacterial and fungal community structure. *Sci Total Environ*. 2024, 949:175127. doi: 10.1016/j.scitotenv.2024.175127.

- Xu B, Hou Y, Yang Z, Fang J, Wu X, Qi J, Li H (2025). Management of saline-alkali sandy soils by amphoteric lignin-based sand fixation. *International Journal of Biological Macromolecules*, 311, Part 3, 144018. <https://doi.org/10.1016/j.ijbiomac.2025.144018>.
- Yadav M (2025). Nanoparticle-facilitated targeted nutrient delivery in plants: Breakthroughs and mechanistic insights. *Plant Nano Biology*, 12, 100156. <https://doi.org/10.1016/j.plana.2025.100156>.
- Yin Y, Cheng GM, Cheng H. Variation of bacterial community diversity and composition in saline-alkali soils reclaimed with flood irrigation and crop cultivation is driven by salinity and edaphic factors. *Sci Total Environ*. 2025, 958:177865. doi: 10.1016/j.scitotenv.2024.177865.
- Zhang N, Liu Q, Chen C, Zhang C, Atakpa EO, Wei X, Shen Q, Jiang L, Tang J, Sun H, Zheng G, Zhou H (2024). Deciphering the performance and mechanisms of glycolipids in regulating crop growth in coastal saline-alkali soils: Perspectives on soil properties and microbial communities. *Applied Soil Ecology*, 201, 105527. <https://doi.org/10.1016/j.apsoil.2024.105527>.
- Zhao X, Grossart HP (2024). Enhancing crop yield and microbial diversity in saline-affected paddy soil through biochar amendment under aquaculture wastewater irrigation. *European Journal of Soil Biology*, 123, 103681. <https://doi.org/10.1016/j.ejsobi.2024.103681>.
- Zhou T, Zhang L, Yang X, Wu Z, Yang Z, Wang J, Chen N, Ren X, Hu S (2025). Prioritizing microbial functions over soil quality for enhanced multifunctionality in saline-sodic soil remediation. *J Environ Manage*. 379:124731. doi: 10.1016/j.jenvman.2025.124731.
- Zhu D, Sun L, Mao L, Li J, Yan B, Li B, Li X (2025). Combined effects of cropping alfalfa (*Medicago sativa* L.) on the soil pore structure, microbial communities and organic carbon fractions in saline soils. *Applied Soil Ecology*, 208, 105993. <https://doi.org/10.1016/j.apsoil.2025.105993>.